

# Statistical Analysis Plan

## ALL-IN-META-BCG-CORONA

Title: Anytime Live and Leading Interim\* meta-analysis of the impact of Bacillus Calmette-Guérin vaccination in health care workers during the SARS-CoV-2 pandemic

Project partners: UMCU / CWI / UvA

SAP version - date: V1.0 of 17 June 2020\*\*

SAP authors:

Coordinating investigator	C.H. (Henri) van Werkhoven Julius Center for Health Sciences and Primary Care University Medical Center Utrecht Heidelberglaan 100, 3584 CX Utrecht, the Netherlands E-mail: c.h.vanwerkhoven@umcutrecht.nl
Trial statistician	Judith ter Schure Machine Learning group Centrum Wiskunde & Informatica (CWI) (the national research institute for mathematics and computer science in the Netherlands) Science Park 123, 1098 XG Amsterdam, the Netherlands E-mail : judith.ter.schure@cw.nl

\* ALL-IN meta-analysis stands for Anytime Live and Leading INterim meta-analysis. This statistical methodology provides meta-analysis of interim data that can be updated at Anytime – so after each new observed data point – Live – so without the need to prespecify when to look – and Leading – because the evidence in the meta-analysis can inform whether individual trials should be stopped or expanded.

This Statistical Analysis Plan proposes to perform ALL-IN meta-analysis for time-to-event data, using the Safe logrank test and Safe hazard ratio confidence sequences. The analysis is literally ALL-IN: any new trial result can be included, even if the decision to start or expand a trial was based on the data already available. For details, see : <https://projects.cwi.nl/safestats/>

\*\* This document received a minor update on 3 July 2020 replacing ‘left-censored’ with ‘left-truncated’ on page 9.

# Signature page

Title: Anytime Live and Leading Interim meta-analysis of the impact of Bacillus Calmette-Guérin vaccination in health care workers during the SARS-CoV-2 pandemic

Document release - date: Statistical analysis plan V1.0 of 17 June 2020

**Approval page** *Signed on 17-06-2020, signatures not available in web version*

<p><b>Henri van Werkhoven</b> <b>Coordinating principle investigator</b> Julius Center for Health Sciences and Primary Care UMC Utrecht</p> <p>Date:</p> <p>Signature:</p>	<p><b>Judith ter Schure</b> <b>Trial statistician</b> Machine Learning group CWI</p> <p>Date:</p> <p>Signature:</p>
--	---

# Table of contents

<b>1</b>	<b>Overview of the study</b> .....	<b>5</b>
<b>2</b>	<b>Definitions</b> .....	<b>6</b>
2.1	Treatment groups .....	6
2.2	Analysed populations.....	6
2.3	Study definitions.....	6
<b>3</b>	<b>General statistical considerations</b> .....	<b>7</b>
3.1	General statistical methodology .....	7
3.2	Handling of missing data .....	7
3.3	Descriptive statistics .....	7
3.4	Inferential statistics .....	7
3.5	Interim analyses and publication of the results .....	7
<b>4</b>	<b>Description of included RCTs</b> .....	<b>8</b>
<b>5</b>	<b>Analysis of the primary endpoints</b> .....	<b>9</b>
5.1	Cumulative incidence of COVID-19 disease .....	9
5.2	Cumulative incidence of hospital admission due to COVID-19 disease .....	9
<b>6</b>	<b>Analysis of the secondary endpoints</b> .....	<b>10</b>
6.1	Number of days of unplanned absenteeism for any reason .....	10
6.2	Number of days of unplanned absenteeism because of COVID-19 disease .. Fout! Bladwijzer niet gedefinieerd.	
<b>7</b>	<b>References</b> .....	<b>11</b>

## List of abbreviations

ALL-IN:	Anytime Live and Leading INterim
BCG:	Bacillus Calmette-Guérin
RCT:	Randomised Controlled Trial
RR:	Relative Risk
HR:	Hazard Ratio

---

# 1 Overview of the study

---

Since the start of the SARS-CoV-2 pandemic, several investigators across the world have set up RCTs of BCG vaccination among health care workers. The hypothesis is that the aspecific immune response to BCG provides indirect protection against COVID-19 or against a severe course of COVID-19. Until today, over 10 trials are ongoing or in preparation. This is great, because if BCG is effective, more trials will ensure that the putative effect will be found, probably earlier and with a more precise estimate of the magnitude of the effect. However, with multiple parallel trials there is an increased risk of false-positive findings, due to multiple testing, which could endanger the completion of the other trials.

If BCG does not prevent COVID-19 (or not to a meaningful extent), each trial needs to limit its chance to incorrectly find superiority: the type-I error or false-positive rate which is normally set at 5%. However, if 10 trials are performed, the chance that at least one of them will be false-positive increases to 40%. Normally, this is not so much of a problem: the majority of trials will not be positive and in the end the conclusion is that there is no effect. However, in this pandemic setting, the collateral damage of a single false-positive finding can be substantial.

We expect that most trials, if not all, will perform interim analyses. The benefit of this is that superiority – if present - will be found earlier, which is important during the pandemic. However, the downside is that, if there is no effect, it is still quite likely that at least one trial will be positive. Since this is found during an interim analysis, this positive trial will most likely be the first to release results. In response to a (possibly false) positive result, investigators of all other trials will need to decide whether or not to stop their trial for ethical reasons and offer BCG to the control group. Also, health care institutions may decide to offer BCG to all health care workers, causing global shortage of BCG. All of this might be in vain if the result acted on is indeed a false-positive. Pooling the evidence across all trials avoids such a scenario. Moreover, prospectively pooling the accruing evidence across all trials has the advantage of increased power, making it likely that a conclusion is reached earlier than in a single trial, when the effect exists.

Normally it will take months to collect and analyse the data of all trials and perform a meta-analysis. In this meta-analysis, central collection of selected data needed for the meta-analysis will already be performed while the trials are ongoing. Data will be analysed on a regular basis using statistical tests and stringent criteria that control false-positive rates. Results will be released as soon as a pooled statistically significant effect is observed or an individual trial yields a statistically significant effect, or when all trials have been completed.

---

## 2 Definitions

---

### 2.1 Treatment groups

BCG: participants randomised to receive BCG

Control: participants randomised to receive no BCG (either no intervention or placebo)

### 2.2 Analysed populations

Participants will be analysed according to the intention-to-treat (ITT) principle, i.e. according to the random allocation.

#### Subgroups:

Participants with documented COVID-19 (time-to-recovery analysis as secondary analysis)

Participants with COVID-19 related hospitalization (time-to-recovery analysis as secondary analysis)

### 2.3 Study definitions

Documented COVID-19 disease is defined as PCR-based detection of SARS-CoV-2 in a respiratory sample.

SARS-CoV-2: severe acute respiratory syndrome coronavirus-2. The virus that causes COVID-19 disease.

---

## 3 General statistical considerations

---

### 3.1 General statistical methodology

The analysis will be performed in the statistical software R.<sup>1</sup> The R-library safestats will be used for most inferential statistics. The version of R and R-libraries used will be reported.

### 3.2 Handling of missing data

Due to the nature of the analysis, missing data are not expected, except in terms of loss-to-follow-up (e.g. withdrawal of participants). Participants with incomplete follow-up will be assumed to have noninformative censoring at the end of their follow-up.

### 3.3 Descriptive statistics

According to the variable type, descriptive statistics will be:

- Quantitative criteria: number of observations (N), mean, standard deviation (SD), median and inter-quartile range (IQR), as applicable.
- Qualitative criteria: number of observations (N), absolute frequency (n) and relative frequency (%). Percentages will be calculated on the number of participants with documented data. Incidence density will be reported (e.g., number of events per 1000 follow-up days) as appropriate.
- Kaplan Meier survival curve for time-to-event endpoints.

### 3.4 Inferential statistics

Safe tests will be used for inferential statistics. Safe tests are designed to control type-I error rates, while maximising flexibility in timing and frequency of interim analyses and allowing for the continuation of data collection beyond the planned trial duration. An explanation of the Safe logrank test for time-to-event data is provided here: <https://projects.cwi.nl/safestats/>.

### 3.5 Interim analyses and publication of the results

Interim analyses of the co-primary endpoint will be performed frequently. Results of the interim analyses will not be published or reported to the trial investigators except in case of one of the following three events:

1. The interim meta-analysis of one or both of the co-primary endpoints yields a statistically significant effect (see section 5 for definition).
2. One of the trials ends with a conclusion. In this case, unveiling the meta-analysis results serves the purpose of confirming or refuting the trial results in order to facilitate an informed decision by all trial investigators and policy makers. This event can happen multiple times as long as trials are ongoing.
3. All trials have been completed and no new trials are being started.

---

## 4 Description of included RCTs

---

We will describe the following characteristics of the included RCTs:

**Meta-data:**

- Country/countries in which the trial has enrolled participants
- Health care setting

**Participant data:**

- Number of participants included and randomised to BCG and control (N)
- Number and percentage of participants with incomplete follow-up for BCG and control (N, %)
- Distribution of actual duration of follow-up per participant for BCG and control (Median, IQR)



---

## 5 Analysis of the primary endpoints

---

Both primary endpoints will be analysed as time-to-event outcome. A Safe logrank test will be used stratified by trial and hospital. The time-scale for the analysis will be calendar time, setting the date of first enrolment of the first trial to  $t = 0$  for all. As a consequence, participants being enrolled later will be left-truncated. This is to ensure that the epidemic wave is on the same time scale for all participants, thus making the hazard function more homogeneous.

### 5.1 Cumulative incidence of COVID-19 disease, weight: 10%

The Safe test for superiority is designed to be a one-sided test for Hazard Ratios (HR) *less than 1* in favour of the treatment, with  $\alpha = 0.25\%$ , and optimised for a minimal clinically relevant HR of 0.8.

The Safe test for harm is designed to be a one-sided test for HR *greater than 1* in favour of the control, with  $\alpha = 0.25\%$ , and optimised for a minimal clinically relevant HR of 1.25.

Results will be reported as a descriptive HR with a 95% two-sided Safe confidence sequence. We will also report a 99.5% two-sided Safe confidence sequence for consistency with the chosen  $\alpha$  for statistical significance.

### 5.2 Cumulative incidence of hospital admission due to COVID-19 disease, weight: 90%

The Safe test for superiority is designed to be a one-sided test for HR *less than 1* in favour of the treatment, with  $\alpha = 2.25\%$ , and optimised for a minimal clinically relevant HR of 0.7.

The Safe test for harm is designed to be a one-sided test for HR *greater than 1* in favour of the control, with  $\alpha = 2.25\%$ , and optimised for a minimal clinically relevant HR of 1.43.

Results will be reported as a descriptive HR with a 95% two-sided Safe confidence sequence. We will also report a 99.5% two-sided Safe confidence sequence for consistency with the chosen  $\alpha$  for statistical significance.

---

## **6 Analysis of the secondary endpoints**

---

The following secondary endpoints will be analysed and reported together with the primary endpoints when results are published:

**6.1 Time-to-recovery from documented COVID-19 disease**

**6.2 Time-to-recovery from hospital admission due to COVID-19 disease**

---

## 7 References

---

1. R Core Team (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
2. <https://projects.cwi.nl/safestats/>